

X-ray Probes of the Universe at Intermediate Redshifts


Ann Hornschemeier
Goddard Space Flight Center





Constellation-X Science Objectives


1




Measure effects of **strong gravity** near the event horizon of black holes.

2

Trace baryonic matter throughout the universe and constrain the nature of **dark matter & dark energy**.



3




Study formation of supermassive **black holes** and trace their evolution with redshift.

- *What roles do they play in galaxy evolution?*
- *What is the total energy output of the universe?*

4

Study the **life cycles of matter** and energy & understand the behavior of matter in extreme environments.

- *What new forms of matter will be discovered?*
- *How does the chemical composition of the universe evolve?*






X-ray Probes at Intermediate Redshift

- **Cosmological studies in the X-ray Band**
 - **Life cycles of matter:**
 - **Accreting binary evolution**
 - **Cosmic energy budget**
 - **Role of BHs in Galaxy Evolution (NO TIME)**
- **Observational requirements**




Intermediate redshift??





Intermediate redshift??



A red rectangular frame with a double-line border, centered on the page. The text "Life Cycles of Matter" is centered within this frame.

Life Cycles of Matter



Life Cycles of Matter

M83, nearby spiral galaxy
with nuclear starburst

Soria & Wu (2003)




- ❑ Accreting binaries → high mass end of IMF, emission persists for very long time (Gigayears)
- ❑ Star formation in heavily obscured areas
- ❑ Supernovae/winds enrich ISM & IGM, affecting star formation and galaxy evolution



Evolutionary Timescales for X-ray Binaries


- ❑ Mass transfer driven by nuclear evolution of donor
- ❑ In high-mass X-ray binaries, timescale is approximately $\sim 2-7 \times 10^6$ yrs (Ghosh & White 2001)
- ❑ low-mass binaries, this timescale is much longer ($> 10^{10}$ yrs)



Credit: <http://chandra.harvard.edu>



X-ray – SFR Correlations




□ **0.5-8 keV luminosity appears to be a reliable SFR indicator**

(Bauer et al. 2002, Seibert, Heckman & Meurer 2002, Ranalli et al. 2002, Cohen et al. 2003)

□ **Dominance of accreting X-ray binaries**



X-ray – SFR Correlations




- More than factor of 3 difference in “star formation coefficients”
- Many X-ray components in galaxies may scale with SFR however (hot ISM, AGN activity?): could differentiate with Con-X spectroscopy



Normal Galaxy X-ray Luminosity Functions

- ❑ Evolution consistent with $(1+z)^3$ but...
- ❑ Conclusions depend on full understanding of L_X -SFR relationship
- ❑ Evolution of binaries?

Norman, Ptak, Hornschemeier et al. 2004






X-rays from Star-forming Galaxies at higher z

- Reddy et al. (2004) find X-ray/SFR relation holds for $1 < z < 3$ galaxies
- Lehmer et al. (2005) have stacked 3000 Lyman break galaxies at $3 < z < 6$, find same conclusions about XR-SFR
- LOCAL analogs to Lyman Breaks: (Heckman et al. 2005)



Steidel et al. 2004






X-ray spectroscopy of post-starburst galaxies

- ❑ Just beyond the reach of current X-ray CCD spectroscopy
- ❑ X-ray binary stellar synthesis models (e.g., Belczynski et al. 2004, Sipior et al. 2005) agree with inferred burst mass fractions and ages in these galaxies (Hornschemeier et al. 2005)
- ❑ May have a tool such as STARBURST99 (Leitherer et al.) for the X-rays by 2010


Goto et al. (2003), Post-starburst Galaxy





X-ray spectroscopic diagnostics of galaxies

SDSS “BPT” Diagram from Brinchmann et al.




A red rectangular frame with a double-line border, featuring small perpendicular tick marks at each of the four corners.

Total Energy Output of the Universe



X-ray Emission from the Universe



Contributors to the
2-10 keV XRB
(Hornschemeier et al. 2003,
Moretti et al. 2003, Bauer et al. 2004)


- ❑ X-ray obscured AGN: 45%
- ❑ X-ray unobscured AGN: 38%
- ❑ Galaxies: 3%
- ❑ Clusters: ~5%

TOTAL: 91%

The high-energy Universe is dominated by accreting supermassive black holes at intermediate redshift



Extragalactic Background Radiation (EBR) Studies



- ❑ Observationally elusive for the longest time: hard X-rays and submm
- ❑ Submm counterparts often optically faint ($I > 24$), VERY difficult to ID with optical spectra
- ❑ X-ray identification rate much better, find AGN aren't dominating submm emission (Alexander et al. 2003)



EBR Studies: Remaining items


- 40 keV background? (HXT!)
- Are Chandra & XMM are missing *highly* obscured AGN? ...might be found in e.g., *Spitzer* surveys (Treister et al. 2005)
- X-rays are most feasible way to identify the physics of the submillimeter sources
→ *constraining >90% of the energy production of the Universe...*



Con-X Spectroscopy of Elusive AGN

- ❑ “Elusive” AGN (e.g., P3, Comastri et al., Fiore et al.)
- ❑ Plausibly an optical aperture effect (Moran et al. 2003, Hornschemeier et al. 2005)
- ❑ May be a class of accreting SMBH that is missed in optical surveys → X-ray spectroscopy alone will reveal the nature of these objects

SDSS Spectrum of
X-ray Luminous Galaxy
(Hornschemeier et al. 2005)






Observational Requirements for Intermediate-Redshift Studies

- ❑ X-ray flux of typical X-ray obscured submm source: $\sim 1 \times 10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$ (0.5-2 keV)
- ❑ X-ray flux of 10^{42} erg/s Seyfert galaxy at $z \sim 1$: $\sim 2 \times 10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$ (0.5-2 keV)
- ❑ X-ray flux of star-forming SDSS galaxy at $z \sim 0.1$: $< 2 \times 10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$
- ❑ Spatial resolution? Not essential at these redshifts (rely on Chandra IDs in some cases)



What if you put an WFI behind this large X-ray mirror?



- ❑ **CONFUSION:** Typical optical magnitudes of AGN at fluxes $< 1 \times 10^{-17} \text{ erg cm}^{-2} \text{ s}^{-1}$ (0.5-2 keV) requires spatial resolution $< 1.6''$
- ❑ Hitting X-ray confusion around $2''$, 5×10^{-18}
- ❑ WFI not so useful for higher-z science at $5''$ - $10''$ planned spatial resolution



Coming soon

- ❑ New science panel to cover this topic, evaluate what is/is not accessible with Con-X
- ❑ Higher-z science available to Con-X but may rely heavily on existing Chandra/XMM surveys
- ❑ Galaxy surveys such as SDSS, GALEX provide excellent samples for projects relevant to life cycles of matter but would be “snapshot” programs to be effective
- ❑ and finally....

Email:
Ann.Hornschemeier.Cardiff@nasa.gov





Coeval Growth of SMBH and Galaxy Bulges

- ❑ **Bulge mass-BH relationship → SFR in spheroids expected to be found around luminous (i.e., *growing*) SMBH**
- ❑ **Based on standard assumptions, the ratio of starburst to AGN emission is a factor of ~5 (Page et al. 2000)**
- ❑ **Initial results by Page et al. 2004**
 - **X-ray unobscured QSOs → not submm emitters**
 - **X-ray obscured QSOs: → strong submm emitters**
 - **Proposes that the largest SMBH are X-ray obscured QSOs for 15% of their lives**